

Math 4441 Practice Midterm 2

November 17, 2022

Name: _____

gtID: _____

Instructions. Read each question carefully and show all your work. Answers without justification will receive little to no credit. Writing your answers in a legible, well-organized manner will maximize your opportunities for partial credit.

This is a closed-note, closed-book exam, and you are expected to abide by the Georgia Tech Honor Challenge. Good luck!

Clearly label any extra papers you want graded.

By signing below, I certify that all work submitted on this exam is my own, and that I have neither given nor received any unauthorized help on this exam.

Question	Points	Score
1	10	
2	10	
3	10	
4	10	
Total:	40	

Formulas

- $df_p = \begin{pmatrix} \frac{\partial f^1}{\partial u^1}(p) & \cdots & \frac{\partial f^1}{\partial u^m}(p) \\ \vdots & \ddots & \vdots \\ \frac{\partial f^n}{\partial u^1}(p) & \cdots & \frac{\partial f^n}{\partial u^m}(p) \end{pmatrix}$
- $d(g \circ f)_p = dg_{f(p)} \cdot df_p$
- $\frac{\partial (g \circ f)^i}{\partial u^j} \Big|_p = \sum_{k=1}^m \frac{\partial g^i}{\partial v^k} \Big|_{f(p)} \frac{\partial f^k}{\partial u^j} \Big|_p$
- $\vec{x}_i := \frac{\partial \vec{x}}{\partial u^i}$
- $\vec{x} = \vec{x} \circ F$ implies $\vec{x}_j = \sum_{i=1}^2 \frac{\partial F^i}{\partial u^j} \vec{x}_i$
- $\vec{n} := \frac{\vec{x}_1 \times \vec{x}_2}{\|\vec{x}_1 \times \vec{x}_2\|}$
- $I_p(\vec{X}, \vec{Y}) := \langle \vec{X}, \vec{Y} \rangle_{\mathbb{R}^3}$
- $g_{ij} = \langle \vec{x}_i, \vec{x}_j \rangle$
- $I_p(\vec{X}, \vec{Y}) = \begin{pmatrix} X^1 \\ X^2 \end{pmatrix}^T \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix} \begin{pmatrix} Y^1 \\ Y^2 \end{pmatrix}$
- $\|\vec{X}\| := \sqrt{I_p(\vec{X}, \vec{X})}$ and $\angle(\vec{X}, \vec{Y}) := \arccos\left(\frac{I_p(\vec{X}, \vec{Y})}{\|\vec{X}\| \cdot \|\vec{Y}\|}\right)$
- $(g^{k\ell}) := (g_{ij})^{-1}$
- $\vec{x} = \vec{x} \circ F$ implies $\tilde{g}_{\alpha\beta} = \sum_{i,j=1}^2 \frac{\partial F^i}{\partial u^\alpha} \frac{\partial F^j}{\partial u^\beta} g_{ij}$
- $\vec{x}_{ij} := \frac{\partial^2 \vec{x}}{\partial u^i \partial u^j}$
- $\kappa_n := \langle \frac{d}{ds} \vec{T}, \vec{n} \rangle$ and $\kappa_g := \langle \frac{d}{ds} \vec{T}, \vec{S} \rangle$
- $L_{ij} := \langle \vec{x}_{ij}, \vec{n} \rangle$
- $\Gamma_{ij}^k := \sum_{\ell=1}^2 \langle \vec{x}_{ij}, \vec{x}_\ell \rangle g^{k\ell}$
- For $\vec{\alpha}$ unit-speed, $\vec{\alpha}'' = \kappa_n \vec{n} + \kappa_g \vec{S}$
- $\kappa_n = \sum_{i,j=1}^2 (\alpha_U^i)' (\alpha_U^j)' L_{ij}$ and $\kappa_g \vec{S} = \sum_{k=1}^2 \left((\alpha_U^k)'' + \sum_{i,j=1}^2 (\alpha_U^i)' (\alpha_U^j)' \Gamma_{ij}^k \right) \vec{x}_k$
- $D_{\vec{v}} f := (f \circ \vec{\alpha})'(0)$, where $\vec{\alpha}(0) = p$ and $\vec{\alpha}'(0) = \vec{v}$
- $\vec{n} = \nu \circ \vec{x}$
- $\mathcal{L}(\vec{v}) := -D_{\vec{v}} \nu$
- $\vec{n}_i := \frac{\partial \vec{n}}{\partial u^i}$
- $\vec{n}_j = -\sum_{i=1}^2 L_j^i \vec{x}_i$
- $(L_{ij}) = (g_{ij})(L_j^i)$, so $(L_j^i) = (g^{k\ell})(L_{ij})$

1. Let $f: U \rightarrow \mathbb{R}$ be a C^∞ function on some open set $U \subset \mathbb{R}^2$. The **graph of f** is the image of the function $\vec{x}: U \rightarrow \mathbb{R}^3$ defined by

$$\vec{x}(u^1, u^2) := (u^1, u^2, f(u^1, u^2)).$$

- (a) (6 points) Verify (using the definition) that \vec{x} is a simple surface.
Hint: You'll need to compute $\vec{x}_1 \times \vec{x}_2$.
- (b) (4 points) Give formulas for the matrices (g_{ij}) and $(g^{k\ell})$ in terms of the function f . You should use the notation $f_i := \frac{\partial f}{\partial u^i}$ for partial derivatives of f , and should simplify your answers.

2. Let $\vec{\alpha}: (a, b) \rightarrow \mathbb{R}^2$ be a unit-speed planar curve. For each $t \in \mathbb{R}$, define a planar curve $\vec{\alpha}_t: (a, b) \rightarrow \mathbb{R}^2$ by

$$\vec{\alpha}_t(s) := \vec{\alpha}(s) + t k(s) \vec{n}(s),$$

where $\vec{n}(s) = J\vec{T}(s)$ is the planar normal vector* for $\vec{\alpha}$ and $k(s)$ is the planar curvature. Note that $\vec{\alpha}_t$ is not necessarily unit-speed. Finally, let $L: \mathbb{R} \rightarrow [0, \infty)$ be defined by

$$L(t) := \ell(\vec{\alpha}_t) = \int_a^b \|\vec{\alpha}'_t(s)\| ds.$$

Note: This problem is way too long to actually show up on a midterm. But you should expect some much smaller version of this type of thing, where you need to know how a technical computation goes.

- (a) (3 points) Prove that

$$L'(t) = \int_a^b \frac{\left\langle \frac{\partial^2}{\partial s \partial t} \vec{\alpha}_t, \frac{\partial}{\partial s} \vec{\alpha}_t \right\rangle}{\sqrt{\left\langle \frac{\partial}{\partial s} \vec{\alpha}_t, \frac{\partial}{\partial s} \vec{\alpha}_t \right\rangle}} ds.$$

- (b) (2 points) Given the formula in part (a), explain why

$$L'(0) = \int_a^b \left\langle \frac{\partial^2}{\partial s \partial t} \vec{\alpha}_t, \frac{\partial}{\partial s} \vec{\alpha}_t \right\rangle \Big|_{t=0} ds.$$

- (c) (3 points) Use integration by parts to show that in fact

$$L'(0) = - \int_a^b \left\langle \frac{\partial}{\partial t} \vec{\alpha}_t, \frac{\partial^2}{\partial s^2} \vec{\alpha}_t \right\rangle \Big|_{t=0} ds.$$

- (d) (2 points) By plugging into the formula just derived, show that

$$L'(0) = \int_a^b -k(s)^2 ds.$$

*In class we just wrote $J\vec{T}$ over and over, but here we're calling it \vec{n} .

3. Consider the simple surface $\vec{x}: (-\pi, \pi) \times \mathbb{R} \rightarrow \mathbb{R}^3$ defined by

$$\vec{x}(u^1, u^2) := (\cos u^1, \sin u^1, u^1 + u^2),$$

whose image is the cylinder $x^2 + y^2 = 1$.

- (a) (4 points) Compute the coefficients L_{ij} of the second fundamental form. Recall that $L_{ij} := \langle \vec{x}_{ij}, \vec{n} \rangle$.
- (b) (6 points) Use the formula $(L_j^i) = (g_{ij})^{-1}(L_{ij})$ to compute the matrix representation (L_j^i) of the Weingarten map.
Note: You will need to compute (g_{ij}) along the way.
- (c) (*Bonus, but not really because there aren't any bonus points, and also this is a practice midterm*)
The matrices $(g_{ij})^{-1}$ and (L_{ij}) are symmetric; how does it happen that (L_j^i) is not symmetric?

4. (a) (3 points) Let $\{\vec{T}(s), \vec{S}(s), \vec{n}(s)\}$ be the Darboux frame of a unit-speed surface curve. Prove that $\frac{d}{ds}\vec{T}(s)$ lies in the plane spanned by $\vec{S}(s)$ and $\vec{n}(s)$.
- (b) (2 points) The following equation[†] tells us why we care about the Christoffel symbols Γ_{ij}^k :

$$\vec{x}_{ij} = L_{ij} \vec{n} + \sum_{k=1}^2 \Gamma_{ij}^k \vec{x}_k.$$

Interpret this equation by writing a sentence that begins with, “The Christoffel symbols tell us...”

- (c) (3 points) Sketch a surface of revolution for which exactly three latitudes are geodesics. Identify these geodesics, and sketch at least two additional geodesics.
- (d) (2 points) Sketch a surface \mathcal{S} and choose a point $p \in \mathcal{S}$ such that the Weingarten map can be represented by the matrix

$$\begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Indicate the point p that you’ve chosen, and **draw the surface normal \vec{n} at this point**. You don’t have to justify your sketch, but you are welcome to add words to explain what you’ve drawn.

[†]Notice that this equation is *not* on your formula sheet. It also won’t be on the formula sheet for the real midterm.